



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW MINOR USE OF A REGISTERED AGVET CHEMICAL
PRODUCT FOR CONTROL OF DOWNY MILDEW IN CANE BERRIES INCLUDING
BLACKBERRIES AND RASPBERRIES**

PERMIT NUMBER – PER93024

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 15 AUGUST 2023 TO 31 AUGUST 2031

Permit Holder:

HORTICULTURE INNOVATION AUSTRALIA LTD
Level 15, 141 Walker Street
NORTH SYDNEY NSW 2060

Persons who can use the product under this permit:

Persons generally

CONDITIONS OF USE

Products to be used:

INFINITO SC FUNGICIDE (APVMA No. 80625)

Containing: 625 g/L PROPAMOCARB HYDROCHLORIDE and 62.5 g/L FLUOPICOLIDE as the only active constituents.

Restraints:

DO NOT apply by aircraft.

DO NOT apply more than three spray applications per season.

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff for at least 3 days after application.

Spray drift restraints

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between three and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a MEDIUM spray droplet size category

DO NOT apply by a vertical sprayer unless the following requirements are met:

- Spray is not directed above the target canopy.
- The outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site.
- For dilute water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas are observed (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for vertical sprayers').

Buffer zones for vertical sprayers

Application rate	Maximum dilute water rate	Type of target canopy	Mandatory downwind buffer zones
160 mL/100 L	1000 L/ha	All	Natural aquatic areas 5 metres

Directions for Use:

Crop	Pest	Rate
Cane berries, including blackberries and raspberries Field and protected grown crops	Downy mildew (<i>Peronospora</i> spp.)	1.6 L/ha

Critical Use Comments:

- Commence treatment when conditions favour disease development – humid or wet conditions.
- Apply by airblast or boom sprayer.
- Apply with a minimum spray volume of 250 L water/ha to ensure thorough and uniform coverage of all plant foliage.
- Apply no more than three (3) applications per crop.
- Apply at 7–10 day intervals. Use the shorter spray interval under heavy disease pressure.
- A non-ionic wetting agent, e.g. Agral 600 should be added at 10 mL/100 L of spray solution.
- Use in accordance with a Fungicide Resistance Management Strategy.
- Follow all label Safety directions and First aid instructions.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

Withholding Period:

Harvest: Do not harvest for 1 DAY after application.

Jurisdiction:

All States and Territories, except VIC.

Note: Victoria is not included in this permit, as their Control-of-Use legislation means a permit is not required to legalise this off-label use in that State.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

Export of treated produce

Temporary Maximum Residue Limits (TMRLs) have been established to allow treated produce to be used for human consumption. TMRL have been established for FLUOPICOLIDE and PROPAMOCARB in CANE BERRIES. MRLs can be found in the *Agricultural and Veterinary Chemicals Code (MRL Standard) Instrument 2023*. MRLs apply only to produce marketed and consumed in Australia. If treated produce is to be exported, residues must not exceed the limits/tolerances of the importing country.

The holder of the permit must notify the APVMA of new information, including relevant information in accordance with section 161 of the Schedule to the *Agricultural and Veterinary Chemicals Code Act 1994*, in accordance with the obligation imposed by that section.

Issued by the Australian Pesticides and Veterinary Medicines Authority

Note: 25/08/2026 – Permit issued as Version 2 with expiry 31/8/2031